

# Independence and Counting



Berlin Chen  
Department of Computer Science & Information Engineering  
National Taiwan Normal University



## Reference:

- D. P. Bertsekas, J. N. Tsitsiklis, *Introduction to Probability* , Sections 1.5-1.6

## Independence (1/2)

- Recall that conditional probability  $\mathbf{P}(A|B)$  captures the partial information that event  $B$  provides about event  $A$
- A special case arises when the occurrence of  $B$  provides no such information and does not alter the probability that  $A$  has occurred

$$\mathbf{P}(A|B) = \mathbf{P}(A)$$

–  $A$  is independent of  $B$  (  $B$  also is independent of  $A$  )

$$\Rightarrow \mathbf{P}(A|B) = \frac{\mathbf{P}(A \cap B)}{\mathbf{P}(B)} = \mathbf{P}(A)$$

$$\Rightarrow \mathbf{P}(A \cap B) = \mathbf{P}(A)\mathbf{P}(B)$$


## Independence (2/2)

- $A$  and  $B$  are independent  $\Rightarrow A$  and  $B$  are disjoint (?)

– No ! Why ?

- $A$  and  $B$  are disjoint then  $P(A \cap B) = 0$
- However, if  $P(A) > 0$  and  $P(B) > 0$

$$\Rightarrow P(A \cap B) \neq P(A)P(B)$$



- Two disjoint events  $A$  and  $B$  with  $P(A) > 0$  and  $P(B) > 0$  are never independent


## Independence: An Example (1/3)

- **Example 1.19.** Consider an experiment involving two successive rolls of a 4-sided die in which all 16 possible outcomes are equally likely and have probability  $1/16$

(a) Are the events,

$$A_i = \{1\text{st roll results in } i\},$$

$B_j = \{2\text{nd roll results in } j\}$ , independent?



Using Discrete Uniform  
Probability Law here

$$\mathbf{P}(A_i \cap B_j) = \frac{1}{16}$$

$$\mathbf{P}(A_i) = \frac{4}{16}, \quad \mathbf{P}(B_i) = \frac{4}{16}$$

$$\Rightarrow \mathbf{P}(A_i \cap B_j) = \mathbf{P}(A_i)\mathbf{P}(B_i)$$

$\Rightarrow A_i$  and  $B_j$  are independent!

## Independence: An Example (2/3)

(b) Are the events,

$$A = \{1\text{st roll is a } 1\},$$

$B = \{\text{sum of the two rolls is a } 5\}$ , independent?

$$\mathbf{P}(A) = \frac{4}{16} \quad (\text{the results of two rolls are } (1,1), (1,2), (1,3), (1,4))$$

$$\mathbf{P}(B) = \frac{4}{16} \quad (\text{the results of two rolls are } (1,4), (2,3), (3,2), (4,1))$$

$$\mathbf{P}(A \cap B) = \frac{1}{16} \quad (\text{the only one result of two rolls is } (1,4))$$

$$\Rightarrow \mathbf{P}(A \cap B) = \mathbf{P}(A)\mathbf{P}(B)$$

$\Rightarrow A$  and  $B$  are independent!

## Independence: An Example (3/3)

(c) Are the events,

$A = \{\text{maximum of the two rolls is } 2\}$ ,

$B = \{\text{minimum of the two rolls is } 2\}$ , independent?

$$P(A) = \frac{3}{16} \quad (\text{the results of two rolls are } (1,2), (2,1), (2,2))$$

$$P(B) = \frac{5}{16} \quad (\text{the results of two rolls are } (2,2), (2,3), (2,4), (3,2), (4,2))$$

$$P(A \cap B) = \frac{1}{16} \quad (\text{the only one result of two rolls is } (2,2))$$

$$\Rightarrow P(A \cap B) \neq P(A)P(B)$$

$\Rightarrow A$  and  $B$  are dependent!

# Conditional Independence (1/2)

- Given an event  $C$ , the events  $A$  and  $B$  are called **conditionally independent** if

$$P(A \cap B | C) = P(A | C)P(B | C) \quad 1$$

- We also know that

$$\begin{aligned} P(A \cap B | C) &= \frac{P(A \cap B \cap C)}{P(C)} \quad \text{multiplication rule} \\ &= \frac{P(C)P(B | C)P(A | B \cap C)}{P(C)} \quad 2 \end{aligned}$$

- If  $P(B | C) > 0$ , we have an alternative way to express conditional independence

$$P(A | B \cap C) = P(A | C) \quad 3$$

## Conditional Independence (2/2)

- Notice that independence of two events  $A$  and  $B$  with respect to the unconditionally probability law does not imply conditional independence , and vice versa

$$\mathbf{P}(A \cap B) = \mathbf{P}(A)\mathbf{P}(B) \quad \cancel{\Leftrightarrow} \quad \mathbf{P}(A \cap B|C) = \mathbf{P}(A|C)\mathbf{P}(B|C)$$

- If  $A$  and  $B$  are independent, the same holds for
  - (i)  $A$  and  $B^c$
  - (ii)  $A^c$  and  $B^c$ 
    - How can we verify it ? (See Problem 38)

# Conditional Independence: Examples (1/2)

- **Example 1.20.** Consider two independent fair coin tosses, in which all four possible outcomes are equally likely. Let

Using Discrete Uniform Probability Law here

$$H_1 = \{\text{1st toss is a head}\}, \quad (H, T), (H, H)$$

$$H_2 = \{\text{2nd toss is a head}\}, \quad (T, H), (H, H)$$

$$D = \{\text{the two tosses have different results}\}. \quad (T, H), (H, T)$$

$$\mathbf{P}(H_1|D) = \frac{1}{2} \quad (H, T)$$

$$\mathbf{P}(H_2|D) = \frac{1}{2} \quad (T, H)$$

$$\mathbf{P}(H_1 \cap H_2|D) = \frac{\mathbf{P}(H_1 \cap H_2 \cap D)}{\mathbf{P}(D)} = 0 \neq \mathbf{P}(H_1|D)\mathbf{P}(H_2|D)$$

$\Rightarrow H_1$  and  $H_2$  are conditionally dependent!

## Conditional Independence: Examples (2/2)

- **Example 1.21.** There are two coins, a blue and a red one
  - We choose one of the two at random, each being chosen with probability 1/2, and proceed with two independent tosses
  - The coins are biased: with the blue coin, the probability of heads in any given toss is 0.99, whereas for the red coin it is 0.01
  - Let  $B$  be the event that the blue coin was selected. Let also  $H_i$  be the event that the  $i$ -th toss resulted in heads

conditional case:  $\mathbf{P}(H_1 \cap H_2 | B) = \mathbf{P}(H_1 | B)\mathbf{P}(H_2 | B)$  Given the choice of a coin, the events  $H_1$  and  $H_2$  are independent

unconditional case:  $\mathbf{P}(H_1 \cap H_2) = ?$   $\mathbf{P}(H_1)\mathbf{P}(H_2)$

$$\mathbf{P}(H_1) = \mathbf{P}(B)\mathbf{P}(H_1 | B) + \mathbf{P}(B^C)\mathbf{P}(H_1 | B^C) = \frac{1}{2} \cdot 0.99 + \frac{1}{2} \cdot 0.01 = \frac{1}{2}$$

$$\mathbf{P}(H_2) = \mathbf{P}(B)\mathbf{P}(H_2 | B) + \mathbf{P}(B^C)\mathbf{P}(H_2 | B^C) = \frac{1}{2} \cdot 0.99 + \frac{1}{2} \cdot 0.01 = \frac{1}{2}$$

$$\mathbf{P}(H_1 \cap H_2) = \mathbf{P}(B)\mathbf{P}(H_1 \cap H_2 | B) + \mathbf{P}(B^C)\mathbf{P}(H_1 \cap H_2 | B^C)$$

$$= \frac{1}{2} \cdot 0.99 \cdot 0.99 + \frac{1}{2} \cdot 0.01 \cdot 0.01 \neq \frac{1}{4}$$

# Independence of a Collection of Events

- We say that the events  $A_1, A_2, \dots, A_n$  are **independent** if

$$\mathbf{P}\left(\bigcap_{i \in S} A_i\right) = \prod_{i \in S} \mathbf{P}(A_i), \text{ for every subset } S \text{ of } \{1, 2, \dots, n\}$$

- For example, the independence of three events  $A_1, A_2, A_3$  amounts to satisfying the four conditions

$$\mathbf{P}(A_1 \cap A_2) = \mathbf{P}(A_1)\mathbf{P}(A_2)$$

$$\mathbf{P}(A_1 \cap A_3) = \mathbf{P}(A_1)\mathbf{P}(A_3)$$

$2^n - n - 1$

$$\mathbf{P}(A_2 \cap A_3) = \mathbf{P}(A_2)\mathbf{P}(A_3)$$

$$\mathbf{P}(A_1 \cap A_2 \cap A_3) = \mathbf{P}(A_1)\mathbf{P}(A_2)\mathbf{P}(A_3)$$

# Independence of a Collection of Events: Examples (1/4)

- **Example 1.22. Pairwise independence does not imply independence.**

- Consider two independent fair coin tosses, and the following events:

$$H_1 = \{ \text{1st toss is a head} \}, \quad (H, T), (H, H)$$

$$H_2 = \{ \text{2nd toss is a head} \}, \quad (T, H), (H, H)$$

$$D = \{ \text{the two tosses have different results} \}. \quad (T, H), (H, T)$$

$$\mathbf{P}(H_1 \cap H_2) = \mathbf{P}(H_1)\mathbf{P}(H_2)$$

$$\mathbf{P}(H_1 \cap D) = \mathbf{P}(H_1)\mathbf{P}(D)$$

$$\mathbf{P}(H_2 \cap D) = \mathbf{P}(H_2)\mathbf{P}(D)$$

$$\text{However, } \mathbf{P}(H_1 \cap H_2 \cap D) = 0 \neq \mathbf{P}(H_1)\mathbf{P}(H_2)\mathbf{P}(D)$$

## Independence of a Collection of Events: Examples (2/4)

- **Example 1.23. The equality**

$$P(A_1 \cap A_2 \cap A_3) = P(A_1)P(A_2)P(A_3)$$

**is not enough for independence.**

- Consider two independent rolls of a fair six-sided die, and the following events:

$$A = \{ \text{1st roll is 1, 2, or 3} \},$$

$$B = \{ \text{1st roll is 3, 4, or 5} \},$$

$$C = \{ \text{the sum of the two rolls is 9} \}.$$

$$P(A \cap B \cap C) = \frac{1}{36} = \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{4}{36} = P(A)P(B)P(C)$$

However,

$$P(A \cap B) = \frac{1}{6} \neq \frac{1}{2} \cdot \frac{1}{2} = P(A)P(B)$$

$$P(A \cap C) = \frac{1}{36} \neq \frac{1}{2} \cdot \frac{4}{36} = P(A)P(C)$$

$$P(B \cap C) = \frac{1}{12} \neq \frac{1}{2} \cdot \frac{4}{36} = P(B)P(C)$$


## Independence of a Collection of Events: Examples (3/4)

- **Example 1.24. Network connectivity.** A computer network connects two nodes  $A$  and  $B$  through intermediate nodes  $C, D, E, F$  ([See next slide](#))
  - For every pair of directly connected nodes, say  $i$  and  $j$ , there is a given probability  $p_{ij}$  that the link from  $i$  to  $j$  is up. We assume that link failures are independent of each other
  - [What is the probability that there is a path connecting  \$A\$  and  \$B\$  in which all links are up?](#)



$$\mathbf{P}(\text{series subsystem succeeds}) = p_1 p_2 \cdots p_n$$

Series Connection




Parallel Connection

$$\begin{aligned}\mathbf{P}(\text{parallel subsystem succeeds}) &= 1 - \mathbf{P}(\text{parallel subsystem fails}) \\ &= 1 - (1 - p_1)(1 - p_2) \cdots (1 - p_n)\end{aligned}$$

## Independence of a Collection of Events: Examples (4/4)

- **Example 1.24. (cont.)**



$$\begin{aligned} \mathbf{P}(C \rightarrow B) &= 1 - (1 - \mathbf{P}(C \rightarrow E \rightarrow B))(1 - \mathbf{P}(C \rightarrow F \rightarrow B)) \\ &= 1 - (1 - 0.8 \cdot 0.9)(1 - 0.95 \cdot 0.85) \\ &= 0.946 \end{aligned}$$

$$\mathbf{P}(A \rightarrow C \rightarrow B) = \mathbf{P}(A \rightarrow C)\mathbf{P}(C \rightarrow B) = 0.9 \cdot 0.946 = 0.851$$

$$\mathbf{P}(A \rightarrow D \rightarrow B) = \mathbf{P}(A \rightarrow D)\mathbf{P}(D \rightarrow B) = 0.75 \cdot 0.95 = 0.712$$

$$\begin{aligned} \therefore \mathbf{P}(A \rightarrow B) &= 1 - (1 - \mathbf{P}(A \rightarrow C \rightarrow B))(1 - \mathbf{P}(A \rightarrow D \rightarrow B)) \\ &= 1 - (1 - 0.851) \cdot (1 - 0.712) = 0.957 \end{aligned}$$

# Recall: Counting in Probability Calculation

- Two applications of the discrete uniform probability law
  - When the sample space  $\Omega$  has a finite number of equally likely outcomes, the probability of any event  $A$  is given by

$$P(A) = \frac{\text{number of elements of } A}{\text{number of elements of } \Omega}$$

- When we want to calculate the probability of an event  $A$  with a finite number of equally likely outcomes, each of which has an already known probability  $p$ . Then the probability of  $A$  is given by

$$P(A) = p \cdot (\text{number of elements of } A)$$


- E.g., the calculation of  $k$  heads in  $n$  coin tosses

# The Counting Principle

- Consider a process that consists of  $r$  stages. Suppose that:
  - (a) There are  $n_1$  possible results for the first stage
  - (b) For every possible result of the first stage, there are  $n_2$  possible results at the second stage
  - (c) More generally, for all possible results of the first  $i - 1$  stages, there are  $n_i$  possible results at the  $i$ -th stage

Then, the total number of possible results of the  $r$ -stage process is

$$n_1 n_2 \cdot \cdot \cdot \cdot n_r$$



# Common Types of Counting

- Permutations of  $n$  objects

$$n! = n \cdot (n - 1) \cdot (n - 2) \cdots 2 \cdot 1$$

- $k$ -permutations of  $n$  objects

$$\frac{n!}{(n - k)!}$$

- Combinations of  $k$  out of  $n$  objects

$$\binom{n}{k} = \frac{n!}{k!(n - k)!}$$

- Partitions of  $n$  objects into  $r$  groups with the  $i$ -th group having  $n_i$  objects

$$\binom{n}{n_1, n_2, \dots, n_r} = \frac{n!}{n_1! n_2! \cdots n_r!}$$

# Summary of Chapter 1 (1/2)

- A probability problem can usually be broken down into a few basic steps:
  1. The description of the sample space, i.e., the set of possible outcomes of a given experiment
  2. The (possibly indirect) specification of the probability law (the probability of each event)
  3. The calculation of probabilities and conditional probabilities of various events of interest.

# Summary of Chapter 1 (2/2)

- Three common methods for calculating probabilities
  - **The counting method:** if the number of outcome is finite and all outcome are equally likely

$$P(A) = \frac{\text{number of elements of } A}{\text{number of elements of } \Omega}$$

- **The sequential method:** the use of the multiplication rule

$$P(\bigcap_{i=1}^n A_i) = P(A_1)P(A_2|A_1)P(A_3|A_1 \cap A_2) \cdots P(A_n|\bigcap_{i=1}^{n-1} A_i)$$

- **The divide-and-conquer method:** the probability of an event is obtained based on a set of conditional probabilities

$$\begin{aligned} P(B) &= P(A_1 \cap B) + \cdots + P(A_n \cap B) \\ &= P(A_1)P(B|A_1) + \cdots + P(A_n)P(B|A_n) \end{aligned}$$

- $A_1, \dots, A_n$  are disjoint events that form a partition of the sample space

# Recitation

- SECTION 1.5 Independence
  - Problems 37, 38, 39, 40, 42